

# DOKTA-CURE 250EW

Tebuconazole 250g/l

A Fungicide for control of Rust in French beans, Yellow Rust and Septoria on Wheat.

**DAKTARI  
WA KUPONYA RUST**



**JOJEMI  
EAST AFRICA LIMITED**

P. O. Box 953 00600 Nairobi,  
T: +254 795 364079 | +254 719 283 963  
E: [jojemieafrica@gmail.com](mailto:jojemieafrica@gmail.com)  
W: [www.jojemieafrica.co.ke](http://www.jojemieafrica.co.ke)

**DESCRIPTION:**

**DOKTA-CURE 250EW** is a Fungicide for control of Rust in French beans, Yellow Rust and Septoria on Wheat.

Crop	Target Disease	Rate	Timing of application
Wheat	Leaf rust, Stem rust, Leaf & Net blotch	0.5-0.75L/ha In 200-300L of Water. 50ml in 20L of water	Apply at the first signs of infection, Before 5% thresh hold of infection is reached. Repeat application after 3-4 weeks if symptoms reappear. NB: Best before head formation
Barley	Barley Leaf rust, Stem rust, Scald, Net blotch, Spot blotch,	0.5-0.75L/ha In 200-300L of Water. 50ml in 20L of water	DOKTA CURE 250EW should be applied in 14 to 21 day intervals for preventative treatments.
French beans, Common beans, Runner beans, Snow peas	Rust, Angular leaf spot, Anthracnose, Powdery Mildew	0.5L/ha In 400-600L of water 10-15ml in 20L of water	Spray when rust infection begins or at budding, whichever is the earlier. Repeat application 10-14 days later. A third application may be necessary when infection occurs early or disease pressure is high.as the first symptoms of infection are noticed. Maximum applications per season:3
Cabbage	Ring spot	0.5L/ha In 400-600L of water 10-15ml in 20L of water	Spray when ringspot infection begins. Repeat application 10-14 days later. A third application may be necessary when infection occurs early or disease pressure is high.
Bananas	Leaf spot (yellow sigatoka), Leaf speckle, black sigatoka	0.5L/ha In 400-600L of water	Regular de-leaving practices to reduce disease inoculum. Old leaves, and leaves with advanced lesions should be removed, or infected parts of the leaf removed, prior to the application of DOKTA CURE 250EW
Pawpaw	Black spot	0.5L/ha In 400-600L of water	Ensure infected plant material is regularly removed and destroyed to reduce inoculum levels. Apply at 14 day intervals. Ensure thorough coverage of leaves and fruit
Peas	Powdery mildew	0.5L/ha In 400-600L of water	Apply at flowering or at first sign of disease, whichever occurs first. A second spray 14 days later may be necessary under favorable conditions.
Rice	Sheath blight, Leaf blast & Neck blast, Glume discoloration (dirty panicle)	0.5L/ha In 400-600L of water	Spray at the first sign of the disease during the tillering stage. A repeat spray 21 to 28 days later may be required. Ensure thorough coverage of stems and leaves.
Coffee	Leaf Rust	0.5L/ha In 400-600L of water	Make the first application as soon as conditions favor the development of the disease. Repeat at 7 to 14 day intervals. Use the higher rate if the crop is dense and disease pressure is high.

**Pre-harvest interval (PHI):** 7 days in French beans.

**AVAILABLE PACKS :** 20ml, 50ml, 100ml, 250ml, 500ml, 1Ltr & 5Ltr

